

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-(6-Hydroxy-1-benzofuran-3-yl)acetamide

D. B. Arunakumar,<sup>a</sup> G. Krishnaswamy,<sup>a</sup> S. Sreenivasa,<sup>a</sup>  
K. J. Pampa,<sup>b</sup> N. K. Lokanath<sup>b</sup> and P. A. Suchetan<sup>c\*</sup><sup>a</sup>Department of Studies and Research in Chemistry, Tumkur University, Tumkur, Karnataka 572 103, India, <sup>b</sup>Department of Studies in Physics, University of Mysore, Manasagangotri, Mysore, India, and <sup>c</sup>Department of Studies and Research in Chemistry, U.C.S., Tumkur University, Tumkur, Karnataka 572 103, India  
Correspondence e-mail: pasuchetan@yahoo.co.in


Received 8 December 2013; accepted 10 December 2013

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.033;  $wR$  factor = 0.081; data-to-parameter ratio = 10.7.

In the title compound,  $\text{C}_{10}\text{H}_9\text{NO}_3$ , the dihedral angle between the benzofuran ring system (r.m.s. deviation for the non-H atoms =  $0.009$  Å) and the  $-\text{C}-\text{C}(\text{O})-\text{N}-$  segment is  $83.76(1)^\circ$ . In the crystal, molecules are linked by  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds, generating (001) sheets, which feature  $C(4)$  and  $C(10)$  chains.

## Related literature

For a related structure and background to benzofurans, see: Arunakumar *et al.* (2014).



## Experimental

## Crystal data

 $\text{C}_{10}\text{H}_9\text{NO}_3$   
 $M_r = 191.18$ Orthorhombic,  $P2_12_12_1$   
 $a = 5.0939(3)$  Å $b = 9.3629(5)$  Å  
 $c = 18.7422(10)$  Å  
 $V = 893.88(9)$  Å<sup>3</sup>  
 $Z = 4$ Cu  $K\alpha$  radiation  
 $\mu = 0.89$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.36 \times 0.29 \times 0.24$  mm

## Data collection

Bruker APEXII CCD  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2009)  
 $T_{\min} = 0.770$ ,  $T_{\max} = 0.808$ 8714 measured reflections  
1459 independent reflections  
1446 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.036$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.081$   
 $S = 1.14$   
1459 reflections  
136 parameters  
H atoms treated by a mixture of  
independent and constrained  
refinement $\Delta\rho_{\max} = 0.15$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.28$  e Å<sup>-3</sup>  
Absolute structure: Flack (1983),  
1927 Friedel pairs  
Absolute structure parameter:  
 $-0.2(2)$ 

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O3}-\text{H}\text{O3}\cdots\text{O1}^{\text{i}}$	0.82	1.92	2.7006 (16)	158
$\text{N1}-\text{H1N}\cdots\text{O3}^{\text{ii}}$	0.92 (3)	2.08 (3)	2.969 (2)	162 (2)
$\text{N1}-\text{H2N}\cdots\text{O1}^{\text{iii}}$	0.92 (3)	2.03 (3)	2.8958 (18)	156 (2)

Symmetry codes: (i)  $x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$ ; (ii)  $x, y - 1, z$ ; (iii)  $x + 1, y, z$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT-Plus* (Bruker, 2009); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the IOE X-ray diffractometer Facility, University of Mysore, Mysore, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB1712).

## References

- Arunakumar, D. B., Nivedita, R. D., Sreenivasa, S., Madan Kumar, S., Lokanath, N. K. & Suchetan, P. A. (2014). *Acta Cryst.* **E70**, o40.  
 Bruker (2009). *APEX2*, *SADABS*, *SAINT-Plus* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2014). E70, o87 [https://doi.org/10.1107/S1600536813033436]

**2-(6-Hydroxy-1-benzofuran-3-yl)acetamide**

**D. B. Arunakumar, G. Krishnaswamy, S. Sreenivasa, K. J. Pampa, N. K. Lokanath and P. A. Suchetan**

**S1. Comment**

As part of our ongoing structural studies of functionalised benzofuran ring systems at the C-2 position (Arunakumar *et al.*, 2014), we now describe the structure of the title compound, (I).

In the title compound, C<sub>10</sub>H<sub>9</sub>NO<sub>3</sub>, the benzofuran ring is almost planar ((r.m.s. deviation for the non-H atoms = 0.009 Å) (Fig. 1). The dihedral angle between the two planes defined by the benzofuran ring and the –C–C(O)–N– segment is 96.24 (1)°. Compared to this, the dihedral angle between the benzofuran ring and the –C–C–N–O segment in the related structure (Arunakumar *et al.*, 2014) is 2.85 (1)°. In the crystal structure, the molecules are linked to one another through strong N1–H1N···O3 and N1–H2N···O1 hydrogen bonds generating C(4) and C(10) chains (Fig. 2). The molecules are further connected into C(10) chains through strong O3–HO3···O1 hydrogen bonds forming helical structure (Fig. 3). Linking of molecules in zig zag pattern through N1–H1N···O3 H-bonds is shown in Fig. 4.


**S2. Experimental**

(6-Hydroxy-1-benzofuran-3-yl)acetic acid (0.015 mmol) was taken in a round bottom flask containing *N,N*-dimethyl formamide (15 ml). To this 1, 1-carbonyldiimidazole (0.023 mmol) was added at 273 K. The reaction mixture was stirred for 45 minutes. Ammonium acetate (0.046 mmol) and triethyl amine (1 ml) were added, the reaction mixture was stirred at room temperature overnight. The completion of the reaction was monitored by thin layer chromatography. After completion of the reaction, the reaction mixture was diluted with ethylacetate and the organic layer was washed with water followed by brine solution. The organic layer was dried over anhydrous sodium sulfate and concentrated to give the crude product. It was further purified by column chromatography by using Ethyl acetate and petroleum ether (8:2) as eluent.

Colourless prisms were obtained from the solvent system: ethyl acetate / methanol (4:1) at room temperature.

**S3. Refinement**

The H atoms were positioned with idealized geometry using a riding model with C–H = 0.93–0.97 Å and O–H = 0.82 Å. The H atoms of the NH<sub>2</sub> group were located in a difference map and later refined freely. The isotropic displacement parameters for all H atoms were set to 1.2 times  $U_{eq}$  (Carbon) and 1.5 times  $U_{eq}$  (Oxygen).



**Figure 1**

Molecular structure of the title compound, showing displacement ellipsoids drawn at the 50% probability level.






Figure 2

Linking of molecules in the crystal structure through N—H $\cdots$ O hydrogen bonds into C(4) and C(10) chains. H-atoms not involved in H-bonding are omitted for clarity.



**Figure 3**

Formation of helical structure through O—H···O hydrogen bonds.

**Figure 4**

Zig-zag pattern observed in the crystal structure.

### 2-(6-Hydroxy-1-benzofuran-3-yl)acetamide

#### Crystal data

$C_{10}H_9NO_3$

$M_r = 191.18$

Orthorhombic,  $P2_12_12_1$

Hall symbol:  $P\ 2ac\ 2ab$

$a = 5.0939\ (3)\ \text{\AA}$

$b = 9.3629\ (5)\ \text{\AA}$

$c = 18.7422\ (10)\ \text{\AA}$

$V = 893.88\ (9)\ \text{\AA}^3$

$Z = 4$

$F(000) = 400$

Prism

$D_x = 1.421\ \text{Mg m}^{-3}$

Melting point: 533 K

Cu  $K\alpha$  radiation,  $\lambda = 1.54178\ \text{\AA}$

Cell parameters from 1234 reflections

$\theta = 4.7\text{--}64.3^\circ$

$\mu = 0.89\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, colourless

$0.36 \times 0.29 \times 0.24\ \text{mm}$

#### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2009)

$T_{\min} = 0.770$ ,  $T_{\max} = 0.808$

8714 measured reflections

1459 independent reflections

1446 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 64.3^\circ$ ,  $\theta_{\min} = 4.7^\circ$

$h = -5 \rightarrow 5$

$k = -10 \rightarrow 10$

$l = -21 \rightarrow 21$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.033$  $wR(F^2) = 0.081$  $S = 1.14$ 

1459 reflections

136 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sitesH atoms treated by a mixture of independent  
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0488P)^2 + 0.1043P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} < 0.001$  $\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$ Absolute structure: Flack (1983), 1927 Friedel  
pairsAbsolute structure parameter:  $-0.2 (2)$ *Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
H2N	0.766 (5)	0.410 (3)	0.6264 (14)	0.072 (7)*
H1N	0.587 (5)	0.324 (3)	0.5742 (14)	0.069 (7)*
C1	0.4600 (3)	1.02627 (16)	0.55047 (8)	0.0361 (4)
C2	0.6218 (3)	0.90706 (18)	0.54019 (9)	0.0404 (4)
H2	0.7494	0.9092	0.5047	0.049*
C3	0.5949 (3)	0.78677 (18)	0.58188 (8)	0.0402 (4)
H3	0.7026	0.7079	0.5747	0.048*
C4	0.4029 (3)	0.78533 (16)	0.63518 (7)	0.0334 (3)
C5	0.2448 (3)	0.90548 (17)	0.64295 (8)	0.0369 (4)
C6	0.2657 (3)	1.02782 (17)	0.60204 (9)	0.0398 (4)
H6	0.1562	1.1061	0.6088	0.048*
C7	0.1208 (4)	0.75102 (19)	0.72418 (8)	0.0452 (4)
H7	0.0293	0.7101	0.7619	0.054*
C8	0.3173 (3)	0.68572 (17)	0.68946 (8)	0.0370 (4)
C9	0.4300 (3)	0.54149 (18)	0.70437 (8)	0.0406 (4)
H9A	0.3482	0.5023	0.7468	0.049*
H9B	0.6166	0.5506	0.7136	0.049*
C10	0.3882 (3)	0.44027 (15)	0.64264 (8)	0.0331 (3)
N1	0.5982 (3)	0.38614 (19)	0.61193 (9)	0.0492 (4)
O1	0.1637 (2)	0.41077 (13)	0.62188 (6)	0.0433 (3)
O2	0.0692 (2)	0.88620 (13)	0.69762 (6)	0.0473 (3)
O3	0.4914 (2)	1.14862 (11)	0.51112 (6)	0.0454 (3)
HO3	0.5821	1.1314	0.4760	0.068*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0386 (8)	0.0371 (8)	0.0326 (7)	−0.0049 (6)	−0.0004 (6)	−0.0042 (6)
C2	0.0363 (9)	0.0486 (9)	0.0364 (7)	−0.0001 (7)	0.0078 (7)	−0.0003 (6)
C3	0.0371 (9)	0.0440 (8)	0.0396 (8)	0.0048 (7)	0.0066 (7)	−0.0023 (6)
C4	0.0306 (8)	0.0402 (8)	0.0294 (7)	−0.0038 (6)	0.0006 (6)	−0.0059 (6)
C5	0.0334 (8)	0.0444 (8)	0.0328 (7)	−0.0052 (7)	0.0050 (6)	−0.0093 (6)
C6	0.0381 (9)	0.0395 (8)	0.0417 (8)	0.0015 (7)	0.0042 (7)	−0.0081 (7)
C7	0.0499 (10)	0.0505 (9)	0.0353 (8)	−0.0090 (8)	0.0103 (7)	−0.0037 (7)
C8	0.0375 (9)	0.0449 (8)	0.0287 (7)	−0.0085 (7)	0.0000 (7)	−0.0056 (6)
C9	0.0411 (9)	0.0495 (9)	0.0312 (7)	−0.0058 (7)	−0.0054 (7)	0.0026 (6)
C10	0.0296 (8)	0.0377 (7)	0.0320 (7)	−0.0027 (6)	−0.0021 (6)	0.0070 (6)
N1	0.0320 (8)	0.0597 (9)	0.0559 (9)	−0.0004 (7)	0.0012 (7)	−0.0097 (8)
O1	0.0290 (6)	0.0567 (7)	0.0444 (6)	−0.0025 (5)	−0.0029 (5)	−0.0082 (5)
O2	0.0476 (7)	0.0508 (7)	0.0434 (6)	−0.0008 (6)	0.0187 (5)	−0.0079 (5)
O3	0.0537 (8)	0.0405 (6)	0.0419 (6)	−0.0003 (5)	0.0081 (5)	0.0008 (5)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—O3	1.3717 (19)	C7—C8	1.341 (2)
C1—C6	1.383 (2)	C7—O2	1.385 (2)
C1—C2	1.401 (2)	C7—H7	0.9300
C2—C3	1.378 (2)	C8—C9	1.494 (2)
C2—H2	0.9300	C9—C10	1.511 (2)
C3—C4	1.398 (2)	C9—H9A	0.9700
C3—H3	0.9300	C9—H9B	0.9700
C4—C5	1.391 (2)	C10—O1	1.2390 (19)
C4—C8	1.447 (2)	C10—N1	1.317 (2)
C5—O2	1.372 (2)	N1—H2N	0.92 (3)
C5—C6	1.382 (2)	N1—H1N	0.92 (3)
C6—H6	0.9300	O3—HO3	0.8200
O3—C1—C6	116.76 (14)	C8—C7—H7	123.7
O3—C1—C2	121.52 (14)	O2—C7—H7	123.7
C6—C1—C2	121.69 (15)	C7—C8—C4	105.80 (15)
C3—C2—C1	120.98 (14)	C7—C8—C9	127.46 (16)
C3—C2—H2	119.5	C4—C8—C9	126.73 (15)
C1—C2—H2	119.5	C8—C9—C10	111.69 (12)
C2—C3—C4	118.87 (15)	C8—C9—H9A	109.3
C2—C3—H3	120.6	C10—C9—H9A	109.3
C4—C3—H3	120.6	C8—C9—H9B	109.3
C5—C4—C3	118.16 (14)	C10—C9—H9B	109.3
C5—C4—C8	105.84 (13)	H9A—C9—H9B	107.9
C3—C4—C8	136.00 (15)	O1—C10—N1	121.80 (15)
O2—C5—C6	124.99 (15)	O1—C10—C9	120.70 (14)
O2—C5—C4	110.44 (14)	N1—C10—C9	117.50 (15)
C6—C5—C4	124.56 (14)	C10—N1—H2N	121.9 (16)

C5—C6—C1	115.73 (15)	C10—N1—H1N	122.0 (17)
C5—C6—H6	122.1	H2N—N1—H1N	116 (2)
C1—C6—H6	122.1	C5—O2—C7	105.37 (13)
C8—C7—O2	112.56 (14)	C1—O3—HO3	109.5
O3—C1—C2—C3	177.50 (15)	O2—C7—C8—C4	−0.22 (19)
C6—C1—C2—C3	−0.7 (3)	O2—C7—C8—C9	178.51 (14)
C1—C2—C3—C4	−0.2 (2)	C5—C4—C8—C7	0.25 (17)
C2—C3—C4—C5	0.9 (2)	C3—C4—C8—C7	179.59 (18)
C2—C3—C4—C8	−178.39 (17)	C5—C4—C8—C9	−178.49 (14)
C3—C4—C5—O2	−179.68 (13)	C3—C4—C8—C9	0.8 (3)
C8—C4—C5—O2	−0.21 (17)	C7—C8—C9—C10	116.36 (19)
C3—C4—C5—C6	−0.8 (2)	C4—C8—C9—C10	−65.2 (2)
C8—C4—C5—C6	178.69 (15)	C8—C9—C10—O1	−59.9 (2)
O2—C5—C6—C1	178.69 (15)	C8—C9—C10—N1	119.67 (16)
C4—C5—C6—C1	−0.1 (2)	C6—C5—O2—C7	−178.81 (16)
O3—C1—C6—C5	−177.48 (14)	C4—C5—O2—C7	0.08 (17)
C2—C1—C6—C5	0.8 (2)	C8—C7—O2—C5	0.09 (18)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O3—HO3 $\cdots$ O1 <sup>i</sup>	0.82	1.92	2.7006 (16)	158
N1—H1N $\cdots$ O3 <sup>ii</sup>	0.92 (3)	2.08 (3)	2.969 (2)	162 (2)
N1—H2N $\cdots$ O1 <sup>iii</sup>	0.92 (3)	2.03 (3)	2.8958 (18)	156 (2)

Symmetry codes: (i)  $x+1/2, -y+3/2, -z+1$ ; (ii)  $x, y-1, z$ ; (iii)  $x+1, y, z$ .